

# The Hilali Conjecture for Hyperelliptic Spaces

Javier Fernández de Bobadilla, Javier Fresán, Vicente Muñoz  
and Aniceto Murillo

**Abstract** The Hilali conjecture predicts that for a simply connected elliptic space, the total dimension of the rational homotopy does not exceed that of the rational homology. Here, we give a proof of this conjecture for a class of elliptic spaces known as hyperelliptic.

**Keywords** Rational homotopy · Sullivan models · Elliptic spaces · Tor functors

## 1 Introduction

Let  $X$  be a simply connected CW-complex. Then,  $X$  is said to be of elliptic type if both  $\dim H^*(X, \mathbb{Q}) < \infty$  and  $\dim \pi_*(X) \otimes \mathbb{Q} < \infty$ . For these spaces, Hilali conjectured in [5] the following:

*Conjecture 1* If  $X$  is a simply connected CW-complex of elliptic type, then

$$\dim \pi_*(X) \otimes \mathbb{Q} \leq \dim H^*(X, \mathbb{Q}).$$

---

J. Fernández de Bobadilla (✉)  
Instituto de Ciencias Matemáticas CSIC-UAM-UC3M-UCM, Consejo Superior de  
Investigaciones Científicas, C/Nicolás Cabrera, nº 13–15, Campus Cantoblanco UAM,  
28049 Madrid, Spain  
e-mail: javier@icmat.es

J. Fresán  
LAGA, UMR 7539, Institut Galilée, Université Paris 13, 99, Avenue Jean-Baptiste Clément,  
93430 Villetaneuse, France  
e-mail: fresan@math.univ-paris13.fr

V. Muñoz  
Facultad de Ciencias Matemáticas, Universidad Complutense de Madrid, Plaza de Ciencias 3,  
28040 Madrid, Spain  
e-mail: vicente.munoz@mat.ucm.es

A. Murillo  
Departamento de Álgebra, Geometría y Topología, Universidad de Málaga, Ap. 59,  
29080 Málaga, Spain  
e-mail: aniceto@uma.es

By the theory of minimal models of Sullivan [3], the rational homotopy type of  $X$  is encoded in a differential algebra  $(A, d)$  called *the minimal model* of  $X$ . This is a free graded algebra  $A = \Lambda V$ , generated by a graded vector space  $V = \bigoplus_{k \geq 2} V^k$ , and with decomposable differential, i.e.,  $d : V^k \rightarrow (\Lambda^{\geq 2} V)^{k+1}$ . It satisfies that:

$$\begin{aligned} V^k &= (\pi_k(X) \otimes \mathbb{Q})^*, \\ H^k(\Lambda V, d) &= H^k(X, \mathbb{Q}). \end{aligned}$$

Therefore, the Hilali conjecture can be rewritten as follows: for a finite-dimensional graded vector space  $V$  (in degrees bigger or equal than two), we have

$$\dim V \leq \dim H^*(\Lambda V, d)$$

for any decomposable differential  $d$  on  $\Lambda V$ .

An elliptic space  $X$  is called of pure type if its minimal model  $(\Lambda V, d)$  satisfies that  $V = V^{even} \oplus V^{odd}$ ,  $d(V^{even}) = 0$  and  $d(V^{odd}) \subset \Lambda V^{even}$ . Also  $X$  is called hyperelliptic if  $d(V^{even}) = 0$  and  $d(V^{odd}) \subset \Lambda^+ V^{even} \otimes \Lambda V^{odd}$ .

In his thesis [5] in 1990, Hilali proved Conjecture 1 for elliptic spaces of pure type. The conjecture is known to hold [6, 7] also in several cases: H-spaces, nilmanifolds, symplectic, and cosymplectic manifolds, coformal spaces with only odd-degree generators, and formal spaces. Hilali and Mamouni [6, 7] have also proved Conjecture 1 for hyperelliptic spaces under various conditions in the homotopical and homological Euler characteristics.

The main result of this paper is the following:

**Theorem 1** *Conjecture 1 holds for hyperelliptic spaces.*

We shall start by proving it for elliptic spaces of pure type in Sect. 3. This requires reducing the question to a problem about Tor functors of certain modules of finite length over a polynomial ring. We solve it by using a semicontinuity result for the Tor functor. Then, in Sect. 4 we prove Theorem 1 for hyperelliptic spaces. For this we have to prove a semicontinuity result for the homology of elliptic spaces, and apply it to reduce the general case to the case in which the minimal model only has generators of odd degree and zero differential. We give two different proofs of an inequality from which the result follows.

## 2 Minimal Models

We recall some definitions and results about minimal models [2]. Let  $(A, d)$  be a *differential algebra*, that is,  $A$  is a (positively) graded commutative algebra over the rational numbers, with a differential  $d$  which is a derivation, i.e.,  $d(a \cdot b) = (da) \cdot b + (-1)^{\deg(a)} a \cdot (db)$ , where  $\deg(a)$  is the degree of  $a$ . We say that  $A$  is connected if  $A^0 = \mathbb{Q}$ , and simply connected if moreover  $A^1 = 0$ .

A simply connected differential algebra  $(A, d)$  is said to be *minimal* if:

1.  $A$  is free as an algebra, that is,  $A$  is the free algebra  $\Lambda V$  over a graded vector space  $V = \bigoplus_{k \geq 2} V^k$ , and
2. For  $x \in V^k$ ,  $dx \in (\Lambda V)^{k+1}$  has no linear term, i.e., it lives in  $\Lambda V^{>0} \cdot \Lambda V^{>0} \subset \Lambda V$ .

Let  $(A, d)$  be a simply connected differential algebra. A minimal model for  $(A, d)$  is a minimal algebra  $(\Lambda V, d)$  together with a quasi-isomorphism  $\rho : (\Lambda V, d) \rightarrow (A, d)$  (that is, a map of differential algebras such that  $\rho_* : H^*(\Lambda V, d) \rightarrow H^*(A, d)$  is an isomorphism). A minimal model for  $(A, d)$  exists and it is unique up to isomorphism.

Now consider a simply connected CW-complex  $X$ . There is an algebra of piecewise polynomial rational differential forms  $(\Omega_{PL}^*(X), d)$  defined in [3, Chap. VIII]. A minimal model of  $X$  is a minimal model  $(\Lambda V, d)$  for  $(\Omega_{PL}^*(X), d)$ . We have that

$$V^k = (\pi_k(X) \otimes \mathbb{Q})^*,$$

$$H^k(\Lambda V, d) = H^k(X, \mathbb{Q}).$$

A space  $X$  is elliptic [1] if both  $\sum \dim \pi_k(X) \otimes \mathbb{Q} < \infty$  and  $\sum \dim H^k(X, \mathbb{Q}) < \infty$ . Equivalently, if  $(\Lambda V, d)$  is the minimal model, we require that both  $V$  and  $H^*(\Lambda V, d)$  are finite dimensional. For elliptic spaces, the Euler–Poincaré and the homotopic characteristics are well defined:

$$\chi = \sum_{i \geq 0} (-1)^i \dim H^i(\Lambda V, \mathbb{Q}),$$

$$\chi_\pi = \sum_{i \geq 0} (-1)^i \dim \pi_i(X) \otimes \mathbb{Q} = \dim V^{even} - \dim V^{odd}.$$

We refer the reader to [2, Theorem 32.10] for the proof of the following:

**Proposition 1** *Let  $(\Lambda V, d)$  be an elliptic minimal model. Then,  $\chi \geq 0$  and  $\chi_\pi \leq 0$ . Moreover,  $\chi_\pi < 0$  if and only if  $\chi = 0$ .*

In his thesis [5], M. Hilali conjectured that for elliptic spaces:

$$\dim \pi_*(X) \otimes \mathbb{Q} \leq \dim H^*(X, \mathbb{Q}).$$

In algebraic terms, this is equivalent to

$$\dim V \leq \dim H^*(\Lambda V, d),$$

whenever  $(\Lambda V, d)$  is a minimal model with  $\dim V < \infty$ . Note that finiteness of both  $\dim H^*(X, \mathbb{Q})$  and  $\dim \pi_*(X) \otimes \mathbb{Q}$  is necessary. Otherwise, one can easily construct counterexamples such as  $X = S^3 \vee S^3$ .

### 3 Proof of the Hilali Conjecture for Elliptic Spaces of Pure Type

A minimal model  $(\Lambda V, d)$  is of *pure type* if  $V = V^{even} \oplus V^{odd}$ , with

$$d(V^{even}) = 0, \quad d(V^{odd}) \subset \Lambda V^{even}.$$

An elliptic space is of pure type if its minimal model is so. These spaces are widely studied in [2, § 32]. By Proposition 1, we have that  $\dim V^{even} - \dim V^{odd} \leq 0$ . Let  $n = \dim V^{even}$  and  $n + r = \dim V^{odd}$ , where  $r \geq 0$ . Write  $x_1, \dots, x_n$  for the generators of even degree, and  $y_1, \dots, y_{n+r}$  for the generators of odd degree. Then,  $dx_i = 0$ , and  $dy_j = P_j(x_1, \dots, x_n)$ , where  $P_j$  are polynomials without linear terms.

In this section, we prove the following:

**Theorem 2** *The Hilali conjecture holds for elliptic spaces of pure type.*

#### 3.1 Expressing the Homology as a Tor Functor

To work over nice modules, we would like to reorder the generators  $y_1, \dots, y_{n+r}$ , so that  $P_1, \dots, P_n$  form a regular sequence in  $\Lambda(x_1, \dots, x_n)$ . Recall that this means that the image of  $P_i$  in  $\Lambda(x_1, \dots, x_n)/(P_1, \dots, P_{i-1})$  is not a zero divisor, for any  $i = 1, \dots, n$ . But this is not possible in general, as shown by the following example.

*Example 1* Let  $V = \mathbb{Q}\langle x_1, x_2, y_1, y_2, y_3 \rangle$ , where  $\deg(x_1) = 2$  and  $\deg(x_2) = 6$ . Define a differential  $d$  on  $\Lambda V$  by

$$dy_1 = x_1^6 + x_2^2, \quad dy_2 = x_1^9 + x_2^3, \quad dy_3 = x_1^4 x_2 + x_1 x_2^2.$$

Then,  $(\Lambda V, d)$  is a pure minimal model. It can be proved that it is elliptic if and only if there exist exact powers of  $x_1$  and  $x_2$ . This is the case, since  $2x_1^{10} = d(x_1^4 y_1 + x_1 y_2 - x_2 y_3)$  and  $2x_2^4 = d(x_2^2 y_1 + x_2 y_2 - x_1^5 y_3)$ . But for the same reason, models  $(\Lambda(x_1, x_2, y_i, y_j), d)$  are not elliptic for any choice of indices  $i, j$ . This amounts to say that  $dy_i, dy_j$  are not a regular sequence in  $\Lambda(x_1, x_2)$ .

However, Halperin showed in [4, Lemma 8] that pure models always admit a basis  $z_1, \dots, z_{n+r}$  of  $V^{odd}$  such that  $dz_1, \dots, dz_n$  is a regular sequence in  $\Lambda(x_1, \dots, x_n)$ . This basis is not necessarily homogeneous but it is possible to preserve the lower grading induced by the number of odd elements, that is

$$(\Lambda V)_q^p = (\Lambda V^{even} \otimes \Lambda^q V^{odd})^p.$$

This grading passes to cohomology and by taking into account the quasi-isomorphisms

$$\begin{aligned} (\Lambda(x_1, \dots, x_n, y_1, \dots, y_{n+r}), d) &\xrightarrow{\sim} (\Lambda(x_1, \dots, x_n, z_1, \dots, z_{n+r}), d) \\ (\Lambda(x_1, \dots, x_n, z_1, \dots, z_n), d) &\xrightarrow{\sim} (\Lambda(x_1, \dots, x_n)/(dz_1, \dots, dz_n), d) \end{aligned}$$

with respect to the lower grading, one deduces that:

$$H_*(\Lambda V, d) \cong H_*(\Lambda(x_1, \dots, x_n)/(dz_1, \dots, dz_n) \otimes \Lambda(z_{n+1}, \dots, z_{n+r}), d).$$

So let  $z_1, \dots, z_{n+r}$  be a basis such that  $dz_1, \dots, dz_n$  form a regular sequence. Put  $P_j = dz_j$  for  $j = 1, \dots, n+r$  and consider the module

$$M = \mathbb{Q}[x_1, \dots, x_n]/(P_1, \dots, P_n)$$

over the ring

$$R = \mathbb{Q}[x_1, \dots, x_n].$$

Consider the ring

$$S = \mathbb{Q}[\lambda_1, \dots, \lambda_r]$$

and the map  $f : S \rightarrow R, \lambda_i \mapsto P_{n+i}$ . Then,  $M$  becomes an  $S$ -module.

Consider also the  $S$ -module

$$\mathbb{Q}_0 = S/(\lambda_1, \dots, \lambda_r).$$

Then, we have the following:

**Proposition 2**  $H_*(\Lambda V, d) \cong \text{Tor}_S^*(M, \mathbb{Q}_0)$ .

*Proof* Let  $U = \langle z_1, \dots, z_n \rangle, W = \langle z_{n+1}, \dots, z_{n+r} \rangle$  so that  $V^{odd} = U \oplus W$ . Then, the map  $(\Lambda V^{even} \oplus U, d) \rightarrow (M, 0)$  is a quasi-isomorphism. Actually, the Koszul complex

$$R \otimes \Lambda^n U \rightarrow R \otimes \Lambda^{n-1} U \rightarrow \dots \rightarrow R \otimes \Lambda^1 U \rightarrow R \rightarrow M$$

is exact, which means that  $(R \otimes \Lambda U, d) \xrightarrow{\sim} (M, 0)$ .

Therefore,

$$(\Lambda V, d) = (R \otimes \Lambda U \otimes \Lambda W, d) \xrightarrow{\sim} (M \otimes \Lambda W, d'), \quad (1)$$

is a quasi-isomorphism, where the differential  $d'$  is defined as zero on  $M$ , and  $d'z_{n+i} = \bar{P}_{n+i} \in M$ . This can be seen as follows: the map (1) is a map of differential algebras. Grading both algebras in such a way that  $\Lambda^k W$  has degree  $k$ , we get two spectral sequences. The map between their  $E_1$ -terms is

$$H^*(R \otimes \Lambda U, d) \otimes \Lambda W \rightarrow M \otimes \Lambda W.$$

As this is a quasi-isomorphism, it follows that the map in the  $E_\infty$ -terms is also an isomorphism. The  $E_\infty$ -terms are the homology of both algebras in (1). So the map (1) is a quasi-isomorphism.

Finally, we have to identify  $H^*(M \otimes \Lambda W, d') \cong \text{Tor}_S^*(M, \mathbb{Q}_0)$ . Note that the homology of  $(M \otimes \Lambda W, d')$  is computed as follows: Take the Koszul complex

$$S \otimes \Lambda^r W \rightarrow S \otimes \Lambda^{r-1} W \rightarrow \dots \rightarrow S \otimes \Lambda^1 W \rightarrow S \rightarrow \mathbb{Q}_0,$$

and tensor it with  $M$  over  $S$  (with the  $S$ -module structure given above), to get

$$(M \otimes_S (S \otimes \Lambda W), d') = (M \otimes \Lambda W, d').$$

The homology of this computes  $\text{Tor}_S^*(M, \mathbb{Q}_0)$ .

**Lemma 1** *Under our assumptions,*

$$\dim \text{Tor}_S^0(M, \mathbb{Q}_0) \geq n + 1 \quad \text{and} \quad \dim \text{Tor}_S^r(M, \mathbb{Q}_0) \geq n + 1.$$

*Proof* Clearly,

$$\text{Tor}_S^0(M, \mathbb{Q}_0) = M \otimes_S \mathbb{Q}_0 = M / (\bar{P}_{n+1}, \dots, \bar{P}_{n+r}) = R / (P_1, \dots, P_{n+r}).$$

As all the polynomials  $P_1, \dots, P_{n+r}$  have no linear part, this module contains the constant and linear monomials at least, so  $\dim \text{Tor}_S^0(M, \mathbb{Q}_0) \geq n + 1$ .

For the other inequality, note that  $\text{Tor}_S^r(M, \mathbb{Q}_0)$  is the kernel of  $M \otimes \Lambda^r W \rightarrow M \otimes \Lambda^{r-1} W$ , i.e., the kernel of

$$(P_{n+1}, \dots, P_{n+r}) : M \rightarrow M \oplus \binom{r}{.} \oplus M. \quad (2)$$

Now we use the following fact: As  $M$  is a complete intersection  $R$ -module (it is the quotient of  $R$  by a regular sequence), it has Poincaré duality in the sense that there is a map  $M \rightarrow \mathbb{Q}$  such that  $\Gamma : M \otimes M \xrightarrow{\text{mult}} M \rightarrow \mathbb{Q}$  is a perfect pairing. Take elements  $\nu, \mu_j \in M$ ,  $j = 1, \dots, n$ , such that

$$\begin{aligned} \Gamma(\nu, x_j) &= 0, \quad j = 1, \dots, n, & \Gamma(\nu, 1) &= 1, \\ \Gamma(\mu_j, x_k) &= \delta_{jk}, \quad j, k = 1, \dots, n, & \Gamma(\mu_j, 1) &= 0, \\ \Gamma(\nu, Q) &= \Gamma(\mu_j, Q) = 0, & & \text{for any quadratic } Q \in R. \end{aligned}$$

Since the elements  $\nu, \mu_j$  are in the kernel of (2) and they are linearly independent, we get  $\dim \text{Tor}_S^r(M, \mathbb{Q}_0) \geq n + 1$ .

### 3.2 Semicontinuity Theorem

We are going to prove a semicontinuity theorem for the Tor functors  $\text{Tor}_S^k(M, \mathbb{Q}_0)$  for flat families of modules  $M$  of finite length (i.e., finite dimensional as  $\mathbb{Q}$ -vector spaces).

Consider a variable  $t$ . A family of  $S$ -modules is a module  $\mathcal{M}$  over  $S[t]$  such that for each  $t_0$ , the  $S$ -module

$$M_{t_0} = \mathcal{M}/(t - t_0)$$

is of finite length. We say that  $\mathcal{M}$  is flat over  $\mathbb{Q}[t]$  if it is a flat  $\mathbb{Q}[t]$ -module, under the inclusion  $\mathbb{Q}[t] \hookrightarrow S[t]$ . Consider  $\mathcal{M}$  as a  $\mathbb{Q}[t]$ -module. Then

$$\mathcal{M} \cong \mathbb{Q}[t]^N \oplus \frac{\mathbb{Q}[t]}{(t - t_1)^{b_1}} \oplus \dots \oplus \frac{\mathbb{Q}[t]}{(t - t_l)^{b_l}},$$

for some  $N \geq 0, l \geq 0, 1 \leq b_1 \leq \dots \leq b_l$ . The module is flat if and only if there is no torsion part, i.e.,  $l = 0$  (to see this, tensor the exact sequence  $0 \rightarrow \mathbb{Q}[t] \xrightarrow{t-t_i} \mathbb{Q}[t] \rightarrow \mathbb{Q}[t]/(t-t_i) \rightarrow 0$  with  $\mathcal{M}$ ). Note that for generic  $\xi$ ,  $\text{length}(M_\xi) = N$ . Therefore, the flatness is equivalent to  $\mathcal{M}/(t-t_i)$  being of length  $N$ , i.e.,

$$\mathcal{M} \text{ is flat} \iff \text{length}(M_t) = N, \forall t.$$

**Lemma 2** For any flat family  $\mathcal{M}$ ,

$$\dim \text{Tor}_S^k(M_0, \mathbb{Q}_0) \geq \dim \text{Tor}_S^k(M_\xi, \mathbb{Q}_0),$$

for generic  $\xi \in \mathbb{Q}$ .

*Proof* Let us resolve  $\mathcal{M}$  as an  $S[t]$ -module:

$$0 \rightarrow S[t]^{a_r} \rightarrow \dots \rightarrow S[t]^{a_0} \rightarrow \mathcal{M} \rightarrow 0. \quad (3)$$

As  $\mathcal{M}$  is flat as  $\mathbb{Q}[t]$ -module, if we tensor the inclusion  $\mathbb{Q}[t] \hookrightarrow S[t]$  by  $\mathcal{M}$  over  $\mathbb{Q}[t]$ , we have that  $\mathcal{M} \hookrightarrow \mathcal{M}$  is an inclusion. Hence, the sequence

$$0 \rightarrow \mathcal{M} \hookrightarrow \mathcal{M} \rightarrow \mathcal{M}/(t) \rightarrow 0$$

is exact. But this sequence is the sequence  $0 \rightarrow S[t] \rightarrow S[t] \rightarrow S[t]/(t) \rightarrow 0$  tensored by  $\mathcal{M}$  over  $S[t]$ . Hence,  $\text{Tor}_{S[t]}^1(\mathcal{M}, S[t]/(t)) = 0$ . Obviously  $\text{Tor}_{S[t]}^j(\mathcal{M}, S[t]/(t)) = 0$  for  $j \geq 2$  (since the resolution  $S[t]/(t)$  has two terms).

Using the above, we can tensor (3)  $\otimes_{S[t]} S[t]/(t)$  to get an exact sequence:

$$0 \rightarrow S^{a_r} \rightarrow \dots \rightarrow S^{a_0} \rightarrow M_0 \rightarrow 0. \quad (4)$$

Now we tensor (4) by  $\otimes_S \mathbb{Q}_0$  and take homology to obtain  $\text{Tor}_S^*(M_0, \mathbb{Q}_0)$ . But

$$(4) \otimes_S \mathbb{Q}_0 = (3) \otimes_{S[t]} \mathbb{Q}_0 = ((3) \otimes_{S[t]} \mathbb{Q}[t]) \otimes_{\mathbb{Q}[t]} \mathbb{Q}[t]/(t) = (5) \otimes_{\mathbb{Q}[t]} \mathbb{Q}[t]/(t),$$

where  $\mathbb{Q}_0 = S[t]/(\lambda_1, \dots, \lambda_r, t)$ , and

$$0 \rightarrow \mathbb{Q}[t]^{a_r} \rightarrow \dots \rightarrow \mathbb{Q}[t]^{a_0} \rightarrow \mathcal{F} = \mathcal{M}/(\lambda_1, \dots, \lambda_r) \rightarrow 0. \quad (5)$$

(This is just a complex, maybe not exact.) Analogously,

$$\mathrm{Tor}_S^*(M_0, \mathbb{Q}_\xi) = H^*((5) \otimes_{\mathbb{Q}[t]} \mathbb{Q}[t]/(t - \xi)).$$

So it remains to see that for a complex  $L_\bullet$  of free  $\mathbb{Q}[t]$ -modules like (5), it holds that

$$\dim H^k(L_\bullet \otimes \mathbb{Q}[t]/(t - \xi)) \leq \dim H^k(L_\bullet \otimes \mathbb{Q}[t]/(t)),$$

for generic  $\xi$ . (Tensor products are over  $\mathbb{Q}[t]$ , which we omit in the notation henceforth.) For proving this, just split (5) as short exact sequences

$$0 \rightarrow Z_i \rightarrow L_i \rightarrow B_{i-1} \rightarrow 0, \quad (6)$$

and note that  $Z_i, B_i$  are free  $\mathbb{Q}[t]$ -modules, being submodules of free modules. So  $Z_i = \mathbb{Q}[t]^{z_i}$  and  $B_i = \mathbb{Q}[t]^{b_i}$ . Now  $0 \rightarrow B_i \rightarrow Z_i \rightarrow H^i(L_\bullet) \rightarrow 0$  gives that

$$H^i(L_\bullet) = \mathbb{Q}[t]^{z_i - b_i} \oplus \text{torsion}.$$

For generic  $\xi$ , we have  $\dim H^i(L_\bullet \otimes \mathbb{Q}[t]/(t - \xi)) = z_i - b_i$ . Hence,

$$\begin{array}{ccccccc} 0 & \rightarrow & Z_i \otimes \mathbb{Q}[t]/(t) & \rightarrow & L_i \otimes \mathbb{Q}[t]/(t) & \rightarrow & B_{i-1} \otimes \mathbb{Q}[t]/(t) & \rightarrow & 0 \\ & & \downarrow & & \parallel & & \downarrow & & \\ 0 & \rightarrow & Z_i(L_\bullet \otimes \mathbb{Q}[t]/(t)) & \rightarrow & L_i \otimes \mathbb{Q}[t]/(t) & \rightarrow & B_{i-1}(L_\bullet \otimes \mathbb{Q}[t]/(t)) & \rightarrow & 0. \end{array}$$

The first sequence is (6) tensored by  $\mathbb{Q}[t]/(t)$ . Thus, the last vertical map is surjective, and the first vertical map is injective.

Therefore, we get

$$\begin{aligned} \dim H^i(L_\bullet \otimes \mathbb{Q}[t]/(t)) &= \dim Z_i(L_\bullet \otimes \mathbb{Q}[t]/(t)) - \dim B_i(L_\bullet \otimes \mathbb{Q}[t]/(t)) \\ &\geq \dim Z_i \otimes \mathbb{Q}[t]/(t) - \dim B_i \otimes \mathbb{Q}[t]/(t) \\ &= \dim H^i(L_\bullet) \otimes \mathbb{Q}[t]/(t) - \dim \mathrm{Tor}_1^{\mathbb{Q}[t]}(H^i(L_\bullet), \mathbb{Q}[t]/(t)) \\ &= z_i - b_i, \end{aligned}$$

where we have used in the third line that there is an exact sequence

$$\begin{aligned} 0 &\rightarrow \mathrm{Tor}_1^{\mathbb{Q}[t]}(H^i(L_\bullet), \mathbb{Q}[t]/(t)) \rightarrow B_i \otimes \mathbb{Q}[t]/(t) \\ &\rightarrow Z_i \otimes \mathbb{Q}[t]/(t) \rightarrow H^i(L_\bullet) \otimes \mathbb{Q}[t]/(t) \rightarrow 0, \end{aligned}$$

and in the fourth line that  $\dim(N \otimes \mathbb{Q}[t]/(t)) = \dim \mathrm{Tor}_1^{\mathbb{Q}[t]}(N, \mathbb{Q}[t]/(t))$  for a torsion  $\mathbb{Q}[t]$ -module  $N$ .



### 3.3 Proof of Theorem 2

We proceed to the proof of the Hilali conjecture for elliptic spaces of pure type. We have to prove that

$$\dim H^*(\Delta V, d) \geq 2n + r.$$

By Proposition 2, we need to prove that  $\dim \text{Tor}_S^*(M, \mathbb{Q}_0) \geq 2n + r$ . Consider the family

$$\mathcal{M} = \frac{\mathbb{Q}[t, x_1, \dots, x_n]}{(P_1 + tx_1, \dots, P_n + tx_n)}.$$

For small  $t$ , the hypersurfaces  $P_1 + tx_1, \dots, P_n + tx_n$  intersect in  $N$  points near the origin accounted with multiplicity, where  $N = \text{length}(M)$ . Therefore,  $\mathcal{M}$  is a flat family. By Lemma 2, it is enough to bound below  $\dim \text{Tor}_S^*(M_\xi, \mathbb{Q}_0)$ . But for generic  $t$ , the hypersurfaces  $P_1 + tx_1, \dots, P_n + tx_n$  intersect in  $N$  distinct points (at least, it is clear that they intersect in several points and the origin is isolated of multiplicity one). Therefore,

$$\text{Tor}_S^k(M_\xi, \mathbb{Q}_0) = \text{Tor}_S^k(\mathbb{Q}_0, \mathbb{Q}_0).$$

This is easily computed to have dimension  $\binom{r}{k}$  (using the Koszul complex). Therefore, using Lemma 1,

$$\begin{aligned} \dim \text{Tor}_S^*(M, \mathbb{Q}_0) &\geq (n + 1) + \sum_{k=1}^{r-1} \dim \text{Tor}_S^k(M, \mathbb{Q}_0) + (n + 1) \\ &\geq 2n + 2 + \sum_{k=1}^{r-1} \dim \text{Tor}_S^k(M_\xi, \mathbb{Q}_0) \\ &= 2n + 2 + \sum_{k=1}^{r-1} \binom{r}{k} = 2n + 2^r \geq 2n + r. \end{aligned}$$

*Remark 1* The above computation works for  $r \geq 1$ . If  $r = 0$  then we have to prove that  $\text{length}(M) \geq 2n$ . But then computing the degree 2 nonzero elements in  $M$ , we have that they are at least  $\binom{n+1}{2} - n$ . So for any  $n$ ,

$$\text{length}(M) \geq 1 + n + \binom{n+1}{2} - n = \frac{1}{2}(n+1)n + 1 \geq 2n.$$

## 4 The Hyperelliptic Case

A minimal model  $(\Delta V, d)$  of elliptic type is *hyperelliptic* if  $V = V^{\text{even}} \oplus V^{\text{odd}}$ , and

$$d(V^{\text{even}}) = 0, \quad d(V^{\text{odd}}) \subset \Lambda^+ V^{\text{even}} \otimes \Delta V^{\text{odd}}. \quad (7)$$

An elliptic space is hyperelliptic if its minimal model is so. Note that elliptic spaces of pure type are in particular hyperelliptic.

By Proposition 1 we have that  $\dim V^{even} - \dim V^{odd} \leq 0$ . Let  $n = \dim V^{even}$  and  $n + r = \dim V^{odd}$ , where  $r \geq 0$ . Write  $x_1, \dots, x_n$  for the generators of even degree, and  $y_1, \dots, y_{n+r}$  for the generators of odd degree. Then,  $dx_i = 0$ , and  $dy_j = P_j(x_1, \dots, x_n, y_1, \dots, y_{j-1})$ , where  $P_j$  do not have linear terms.

In this section, we prove the following:

**Theorem 3** *The Hilali conjecture holds for hyperelliptic spaces.*

## 4.1 Semicontinuity for Elliptic Minimal Models

**Lemma 3** *Let  $V$  be a graded rational finite-dimensional vector space, and let  $d$  be a differential for  $\Delta V \otimes \mathbb{Q}[t]$  such that  $dt = 0$ , where  $t$  has degree 0. Take a non-countable field  $\mathbf{k} \supset \mathbb{Q}$ ,  $V_{\mathbf{k}} = V \otimes \mathbf{k}$ . We denote by  $d_{\xi}$  the differential induced on  $\Delta V_{\mathbf{k}} = \Delta V \otimes \mathbf{k}[t]/(t - \xi)$ , for  $\xi \in \mathbf{k}$ . Then*

$$\dim H(\Delta V_{\mathbf{k}}, d_{\xi}) \leq \dim H(\Delta V, d_0),$$

for generic  $\xi \in \mathbf{k}$ .

*Proof* Write

$$0 \rightarrow \tilde{K} \rightarrow \Delta V \otimes \mathbf{k}[t] \rightarrow \tilde{I} \rightarrow 0,$$

where  $\tilde{K}$  and  $\tilde{I}$  are the kernel and image of  $d$ , respectively. Note that both  $\tilde{K}$  and  $\tilde{I}$  are free  $\mathbf{k}[t]$ -modules, being submodules of  $\Delta V \otimes \mathbf{k}[t]$ .

Denote by  $\mathbf{k}_{\xi} = \mathbf{k}[t]/(t - \xi)$ . Then, we have a diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \tilde{K} \otimes \mathbf{k}_{\xi} & \rightarrow & (\Delta V \otimes \mathbf{k}[t]) \otimes \mathbf{k}_{\xi} & \rightarrow & \tilde{I} \otimes \mathbf{k}_{\xi} \rightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \rightarrow & K & \rightarrow & \Delta V_{\mathbf{k}} & \rightarrow & I \rightarrow 0. \end{array} \quad (8)$$

(Here, the tensor products of all  $\mathbf{k}[t]$ -modules are over  $\mathbf{k}[t]$ , and the tensor product  $\Delta V \otimes \mathbf{k}[t]$  is over the rationals.) Therefore, the last vertical map is a surjection, and the first map is an injection.

We have

$$0 \rightarrow \tilde{I} \rightarrow \tilde{K} \rightarrow H(\Delta V \otimes \mathbf{k}[t], d) \rightarrow 0,$$

which is an exact sequence of  $\mathbf{k}[t]$ -modules. Then,  $H(\Delta V \otimes \mathbf{k}[t], d)$  contains a free part and a torsion part. The torsion is supported at some points, which are at most countably many. Therefore, for generic  $\xi \in \mathbf{k}$ ,

$$0 \rightarrow \tilde{I} \otimes \mathbf{k}_{\xi} \rightarrow \tilde{K} \otimes \mathbf{k}_{\xi} \rightarrow H(\Delta V \otimes \mathbf{k}[t], d) \otimes \mathbf{k}_{\xi} \rightarrow 0$$

is exact. As  $\tilde{I} \otimes \mathbf{k}_{\xi} \rightarrow I \subset K$  and  $\tilde{I} \otimes \mathbf{k}_{\xi} \subset \tilde{K} \otimes \mathbf{k}_{\xi} \subset K$ , we have that the last map in (8) is an injection, therefore an isomorphism, and thus the first map is also an isomorphism by the snake lemma.

Note that also, when tensoring with  $\mathbf{k}(t)$ , we have an exact sequence

$$0 \rightarrow \tilde{I} \otimes \mathbf{k}(t) \rightarrow \tilde{K} \otimes \mathbf{k}(t) \rightarrow H(\Delta V \otimes \mathbf{k}[t], d) \otimes \mathbf{k}(t) \rightarrow 0.$$

Also  $H(\Delta V \otimes \mathbf{k}[t], d) \otimes \mathbf{k}(t) = H(\Delta V \otimes \mathbf{k}(t), d)$ , since  $\mathbf{k}(t)$  is a flat  $\mathbf{k}[t]$ -module. Hence,

$$\begin{aligned} \dim H(\Delta V_{\mathbf{k}}, d_{\xi}) &= \dim K - \dim I \\ &= \dim \tilde{K} \otimes \mathbf{k}_{\xi} - \dim \tilde{I} \otimes \mathbf{k}_{\xi} \\ &= \dim H(\Delta V \otimes \mathbf{k}(t), d). \end{aligned}$$

In the first line, we mean  $\dim K - \dim I = \sum_{d \geq 0} (\dim K^d - \dim I^d)$ .

Take now  $\xi = 0$ . The map  $\tilde{K} \rightarrow K \rightarrow K/I$  factors as  $\tilde{K}/\tilde{I} \rightarrow K/I$ . Tensor this map by  $\mathbf{k}_0$  to get  $(\tilde{K}/\tilde{I}) \otimes \mathbf{k}_0 \rightarrow K/I$ . Note that there is an exact sequence

$$\tilde{I} \otimes \mathbf{k}_0 \rightarrow \tilde{K} \otimes \mathbf{k}_0 \rightarrow (\tilde{K}/\tilde{I}) \otimes \mathbf{k}_0 \rightarrow 0,$$

but the first map may not be injective. Then, there is a map

$$\frac{\tilde{K} \otimes \mathbf{k}_0}{\text{Im}(\tilde{I} \otimes \mathbf{k}_0)} = (\tilde{K}/\tilde{I}) \otimes \mathbf{k}_0 \rightarrow K/I.$$

By (8), this is an inclusion. Now we have:

$$\begin{aligned} \dim H(\Delta V, d_{\xi}) &= \dim H(\Delta V \otimes \mathbf{k}(t), d) \\ &= \dim (\tilde{K}/\tilde{I}) \otimes \mathbf{k}(t) \\ &\leq \dim (\tilde{K}/\tilde{I}) \otimes \mathbf{k}_0 \\ &= \dim \frac{\tilde{K} \otimes \mathbf{k}_0}{\text{Im}(\tilde{I} \otimes \mathbf{k}_0)} \\ &\leq \dim K/I \\ &= \dim H(\Delta V_{\mathbf{k}}, d_0) \\ &= \dim_{\mathbb{Q}} H(\Delta V, d_0). \end{aligned}$$

## 4.2 Perturbing the Minimal Model

Let  $x_1, \dots, x_n$  denote generators for  $V^{\text{even}}$ , and  $y_1, \dots, y_{n+r}$  generators for  $V^{\text{odd}}$ . Here,  $dx_i = 0$  and  $dy_j = P_j(x_1, \dots, x_n, y_1, \dots, y_{j-1})$ .

We consider the algebra

$$(\Lambda W, d) = (\Lambda V, d) \otimes (\Lambda \bar{y}_1, 0),$$

where  $\deg(\bar{y}_1) = \deg(x_1) - 1$ . Then

$$\dim H(\Lambda W, d) = 2 \dim H(\Lambda V, d).$$

Consider now the differential  $\delta$  on  $\Lambda W$  such that  $\delta x_j = 0$ ,  $\delta y_j = 0$  and  $\delta \bar{y}_1 = x_1$ . Hence,  $\delta^2 = 0$  and  $d\delta = \delta d = 0$ . So,

$$d_t = d + t\delta$$

is a differential on  $\Lambda W \otimes \mathbf{k}[t]$ .

For generic  $\xi \in \mathbf{k}$ ,  $(\Lambda W_{\mathbf{k}}, d_{\xi})$  verifies that  $d_{\xi} \bar{y}_1 = \xi x_1$ . So, for nonzero  $\xi$ , there is a KS-extension [8, § 1.4]

$$(\Lambda(x_1, \bar{y}_1), d_{\xi}) \longrightarrow (\Lambda W_{\mathbf{k}}, d_{\xi}) \longrightarrow (\Lambda(x_2, \dots, x_n, y_1, \dots, y_{n+r}), d).$$

As  $H(\Lambda(x_1, \bar{y}_1), d_{\xi}) = \mathbf{k}$ , we have that

$$H(\Lambda W_{\mathbf{k}}, d_{\xi}) \cong H(\Lambda(x_2, \dots, x_n, y_1, \dots, y_{n+r}), d).$$

Now we apply Lemma 3 to this to obtain that

$$\dim H(\Lambda(x_2, \dots, x_n, y_1, \dots, y_{n+r}), d) \leq \dim H(\Lambda W, d) = 2 \dim H(\Lambda V, d).$$

Repeating the argument  $n$  times, we get that

$$\dim H(\Lambda(y_1, \dots, y_{n+r}), d) \leq 2^n \dim H(\Lambda V, d).$$

But the hyperelliptic condition says that  $d = 0$  for the first space, so

$$2^n \dim H(\Lambda V, d) \geq \dim H(\Lambda(y_1, \dots, y_{n+r}), d) = 2^{n+r}.$$

This gives

$$\dim H(\Lambda V, d) \geq 2^r. \tag{9}$$

### 4.3 Another Proof of Inequality (9)

In this section, we present a different proof of the inequality  $\dim H(\Lambda V, d) \geq 2^r$  for hyperelliptic spaces. Recall that if  $A$  is a commutative graded differential algebra, and if  $M, N$  are differential graded  $A$ -modules, the differential Tor is defined as

$$\mathrm{Tor}^*(M, N) = H^*(P \otimes_A N),$$

where  $P \xrightarrow{\sim} M$  is a semifree resolution, i.e., a quasi-isomorphism from a semifree  $A$ -module  $P$  to  $M$  (see [2, § 6]).

**Lemma 4** *Let  $C \xleftarrow{\varphi} A \xrightarrow{\psi} B$  be morphisms of commutative differential graded algebras. There exists a convergent spectral sequence*

$$E_2^{p,q} = H^p(B) \otimes \mathrm{Tor}_A^q(\mathbb{Q}, C) \Rightarrow \mathrm{Tor}_A^{p+q}(B, C).$$

*Proof* Decompose  $\varphi$  and  $\psi$  as

$$\begin{array}{ccc} A & \longrightarrow & A \otimes \Lambda W \\ & \searrow \psi & \downarrow \alpha \sim \\ & & B \end{array} \qquad \begin{array}{ccc} A & \longrightarrow & A \otimes \Lambda U \\ & \searrow \varphi & \downarrow \beta \sim \\ & & C \end{array}$$

Then,  $\alpha : A \otimes \Lambda W \xrightarrow{\sim} B$  is a semifree resolution of  $B$  regarded as  $A$ -module, so

$$\mathrm{Tor}_A^*(B, C) = H^*((A \otimes \Lambda W) \otimes_A C).$$

Moreover,  $\mathrm{Id} \otimes \beta : (A \otimes \Lambda W) \otimes_A A \otimes \Lambda U \xrightarrow{\sim} (A \otimes \Lambda W) \otimes_A C$  is a quasi-isomorphism and  $(A \otimes \Lambda W) \otimes_A (A \otimes \Lambda U) \cong A \otimes \Lambda W \otimes \Lambda U$ . Therefore, one gets a rational fibration

$$A \otimes \Lambda W \rightarrow A \otimes \Lambda W \otimes \Lambda U \rightarrow \Lambda U,$$

whose associated Serre spectral sequence has the form

$$E_2^{p,q} = H^p(A \otimes \Lambda W) \otimes H^q(\Lambda U) \Rightarrow H^{p+q}(A \otimes \Lambda W \otimes \Lambda U).$$

On the one hand,  $H^*(A \otimes \Lambda W) = H^*(B)$ . On the other hand, since  $\beta$  is a semifree resolution of  $C$ , we have that

$$H^*(\Lambda U) = H^*((A \otimes \Lambda U) \otimes_A \mathbb{Q}) = \mathrm{Tor}_A^*(\mathbb{Q}, C).$$

Putting all pieces together, we get

$$E_2^{p,q} = H^p(B) \otimes \mathrm{Tor}_A^q(\mathbb{Q}, C) \Rightarrow \mathrm{Tor}_A^{p+q}(B, C).$$

**Theorem 4** *Let  $(\Lambda V, d)$  be a hyperelliptic minimal model. Then*

$$\dim H(\Lambda V, d) \geq 2r.$$

*Proof* Write as usual  $x_1, \dots, x_n$  for generators of  $X = V^{\mathrm{even}}$  and  $y_1, \dots, y_{n+r}$  for generators of  $Y = V^{\mathrm{odd}}$ . When we apply the previous lemma to morphisms  $\mathbb{Q} \leftarrow \Lambda X \hookrightarrow \Lambda V$  we get a spectral sequence

$$E_2 = H(\Lambda V, d) \otimes \mathrm{Tor}_{\Lambda X}^*(\mathbb{Q}, \mathbb{Q}) \Rightarrow \mathrm{Tor}_{\Lambda X}^*(\Lambda V, \mathbb{Q}).$$

On the one hand,

$$\mathrm{Tor}_{\Lambda X}^*(\mathbb{Q}, \mathbb{Q}) = H^*(\Lambda(\bar{x}_1, \dots, \bar{x}_n), 0) = \Lambda(\bar{x}_1, \dots, \bar{x}_n),$$

where  $\Lambda(x_1, \dots, x_n, \bar{x}_1, \dots, \bar{x}_n) \xrightarrow{\sim} \mathbb{Q}$  is a semifree resolution of  $\mathbb{Q}$  regarded as  $\Lambda X$ -module. Hence,  $\bar{x}_i$  are all of odd degree.

On the other hand,  $\Lambda V$  is already  $\Lambda X$ -semifree, so

$$\mathrm{Tor}_{\Lambda X}^*(\Lambda V, \mathbb{Q}) = H(\Lambda V \otimes_{\Lambda X} \mathbb{Q}) = H^*(\Lambda(y_1, \dots, y_{n+k}), 0) = \Lambda(y_1, \dots, y_{n+k}).$$

Then, the inequality

$$\dim H^*(\Lambda V, d) \cdot \dim \mathrm{Tor}_{\Lambda X}^*(\mathbb{Q}, \mathbb{Q}) \geq \dim \mathrm{Tor}_{\Lambda X}^*(\Lambda V, \mathbb{Q})$$

coming from the spectral sequence translates into

$$2^n \dim H^*(\Lambda V, d) \geq 2^{n+r},$$

so the result follows.

#### 4.4 Proof of Theorem 3

Now we prove the inequality  $\dim H(\Lambda V, d) \geq 2n + r$ , for the hyperelliptic minimal model.

If  $r = 0$ , then  $\chi_\pi = 0$ . So [2, Prop. 32.10] says that the model is pure, and this case is already covered by Remark 1.

If  $r > 0$ , then  $\chi_\pi < 0$ . So by Proposition 1,  $\chi = 0$ , and hence, it is enough to prove that

$$\dim H^{even}(\Lambda V, d) \geq n + \frac{r}{2}.$$

Suppose that  $r = 1, 2$ . As the degree 0 and degree 1 elements give always nontrivial homology classes, then  $\dim H^{even}(\Lambda V, d) \geq n + 1$ , and we are done.

Therefore, we can assume  $r \geq 3$ . We use the following fact: if  $P(x)$  is a quadratic polynomial on the  $x$ , and  $P(x) = d\alpha$ ,  $\alpha \in \Lambda V$ , then  $\alpha$  must be linear,  $\alpha \in V^{odd}$  and denoting by  $d_o$  the composition

$$V^{odd} \longrightarrow \Lambda^+ V^{even} \otimes \Lambda V^{odd} \twoheadrightarrow \Lambda^+ V^{even},$$

we have  $P(x) = d_o \alpha$ . So there are at least  $\binom{n+1}{2} - (n+r)$  quadratic terms in the homology. Conjecture 1 is proved if

$$\begin{cases} \text{either} & 1 + n + \binom{n+1}{2} - (n+r) \geq n + \frac{r}{2}, \\ \text{or} & 2^r \geq 2n + r. \end{cases} \quad (10)$$

So now assume that (10) does not hold. Then

$$2^r - r \leq 2n - 1, \quad (11)$$

and  $1 + \binom{n+1}{2} - n < \frac{3}{2}r$ , i.e.,

$$(2n - 1)^2 \leq 12r - 11. \quad (12)$$

Putting together (11) and (12), we get  $2r - r \leq \sqrt{12r - 11}$ , i.e.,  $2r \leq r + \sqrt{12r - 11}$ . This is easily seen to imply that  $r \leq 3$ . So,  $r = 3$  and  $n = 3$ .

It remains to deal with the case  $n = 3$ ,  $r = 3$ , and  $d_o$  is an isomorphism of the odd degree elements onto  $\Lambda^2 V^{even}$ . Let  $x_1, x_2, x_3$  be the even-degree generators, of degrees  $d_1 \leq d_2 \leq d_3$ , respectively. The degrees of  $x_1^2, x_1x_2, x_2^2, x_1x_3, x_2x_3, x_3^2$  are the six numbers

$$2d_1 \leq d_1 + d_2 \leq 2d_2, \quad d_1 + d_3 \leq d_2 + d_3 \leq 2d_3.$$

We have two cases:

- Case  $2d_2 \leq d_1 + d_3$ . We can arrange the odd generators  $y_1, \dots, y_6$  with increasing degree and so that  $d_o y_1 = x_1^2, d_o y_2 = x_1x_2, d_o y_3 = x_2^2, d_o y_4 = x_1x_3, d_o y_5 = x_2x_3, d_o y_6 = x_3^2$ . Clearly,  $dy_1 = x_1^2$ . Then,  $dy_2 = x_1x_2 + P(x_1)$ , where  $P(x_1)$  is a polynomial on  $x_1$ , i.e., of the form  $c x_1^n$ ,  $n \geq 2$ . But this can be absorbed by a change of variables  $y_2 \mapsto y_2 - c x_1^{n-2} y_1$ . So, we can write  $dy_2 = x_1x_2$ . Now the even-degree closed elements in  $\Lambda(x_1, x_2, x_3, y_1, y_2)$  are again polynomials on  $x_1, x_2, x_3$ . So, we can assume  $dy_3 = x_2^2$  as before. Continuing the computation, the even-degree closed elements in  $\Lambda(x_1, x_2, x_3, y_1, y_2, y_3)$  are either polynomials on the  $x_i$ 's or a multiple of the element  $x_2^2 y_1 y_2 - x_1 x_2 y_1 y_3 + x_1^2 y_2 y_3 = d(y_1 y_2 y_3)$ , which is exact. Therefore, we can again manage to arrange that  $dy_4 = x_1x_3$ .
- Case  $2d_2 > d_1 + d_3$ . Then, we have that  $d_o y_3 = x_1x_3$  and  $d_o y_4 = x_2^2$ . As before, we can arrange  $dy_3 = x_1x_3$ . Now the even-degree closed elements in  $\Lambda(x_1, x_2, x_3, y_1, y_2, y_3)$  are polynomials on the  $x_i$ 's or a multiple of  $x_3 y_1 y_2 - x_2 y_1 y_3 + x_1 y_2 y_3$ . But this element has degree  $3d_1 + d_2 + d_3 - 2 > 2d_2$ , so it must be  $dy_4 = x_2^2$ .

In either case,  $dy_1, dy_2, dy_3, dy_4$  are  $x_1^2, x_1x_2, x_2^2$  and  $x_1x_3$ . Let us assume that we are in the first case to carry over the notation.

Now we compute the even-degree closed elements in  $\Lambda(x_1, x_2, x_3, y_1, y_2, y_3, y_4)$ . These are polynomials on  $x_i$ 's or combinations of

$$\begin{aligned} x_2^2 y_1 y_2 - x_1 x_2 y_1 y_3 + x_1^2 y_2 y_3 &= d(y_1 y_2 y_3), \\ x_3 y_1 y_2 - x_2 y_1 y_4 + x_1 y_2 y_4, \\ x_1 x_3 y_2 y_3 - x_2^2 y_2 y_4 + x_1 x_2 y_3 y_4 &= d(y_2 y_3 y_4), \text{ and} \\ x_1 x_3 y_1 y_3 + x_1^2 y_3 y_4 - x_2^2 y_1 y_4 &= d(y_1 y_3 y_4). \end{aligned}$$

Only the second one is nonexact, but its degree is strictly bigger than  $d_2 + d_3$ . So again we can arrange that  $dy_5 = x_2x_3$ .

Finally, the minimal model is:

$$\begin{cases} dy_1 = x_1^2, \\ dy_2 = x_1x_2, \\ dy_3 = x_2^2, \\ dy_4 = x_1x_3, \\ dy_5 = x_2x_3, \\ dy_6 = x_3^2 + P(x_i, y_j). \end{cases}$$

The even-degree closed elements in  $\Lambda(x_1, x_2, x_3, y_1, y_2, y_3, y_4, y_5)$  contain at least

$$\alpha_1 = x_3 y_2 y_3 + x_1 y_3 y_5 - x_2 y_2 y_5 ,$$

$$\alpha_2 = x_3 y_1 y_2 - x_2 y_1 y_4 + x_1 y_2 y_4 .$$

At most, one of them does not survive in  $H(\Lambda V, d)$ , so proving the existence of at least another even-degree cohomology class. Hence,  $\dim H(\Lambda V, d) \geq 10 \geq 9$ , as required.

## References

1. Félix, Y.: La Dichotomie Elliptique-Hyperbolique en Homotopie Rationnelle, Astérisque 176. Société mathématique de France Paris (1989)
2. Félix, Y., Halperin, S., Thomas, J.-C.: Rational Homotopy Theory. Graduate Texts in Mathematics, vol. 205. Springer, Berlin (2000)
3. Griffiths, P.A., Morgan, J.W.: Rational Homotopy Theory and Differential Forms. Progress in Mathematics, vol. 16. Birkhäuser, Basel (1981)
4. Halperin, S.: Finiteness in the minimal models of Sullivan. Trans. Am. Math. Soc. **230**, 173–199 (1977)
5. Hilali, M.R.: Action du tore  $\mathbb{T}^n$  sur les espaces simplement connexes. Ph.D. thesis, Université catholique de Louvain, Belgique (1990)
6. Hilali, M.R., Mamouni, M.I.: A conjectured lower bound for the cohomological dimension of elliptic spaces. J. Homotopy Relat. Struct. **3**, 379–384 (2008)
7. Hilali, M.R., Mamouni, M.I.: A lower bound of cohomologic dimension for an elliptic space. Topol. Appl. **156**, 274–283 (2008)
8. Oprea, J., Tralle, A.: Symplectic Manifolds with no Kähler Structure. Lecture Notes in Mathematics, vol. 1661. Springer, Berlin (1997)